

Lecture 1: Integrable probabilities



Alexey Bufetov and Vadim Gorin

January 24, 2017

- Lecture 1: Uniformly random domino tilings of the Aztec diamond
- Lectures 1-2: Algebra of symmetric functions; Schur processes.
- Lectures 2-3: RSK-algorithm; Last Passage Percolation.
- Lecture 3: Stochastic six vertex model; Hall-Littlewood processes.
- Lectures 4-5: Global asymptotics of stochastic particle systems: robust results.
- Lectures 6-7: Determinantal processes; correlation functions for Schur processes; local asymptotic limit behavior.
- Lecture 8: Asymptotics of the stochastic six vertex model.


Domino tilings of Aztec diamond

Aztec diamond of size 3.




- Aztec diamond of size n is all lattice squares which are (fully) contained in $\{(x, y) : |x| + |y| \leq n + 1\}$.
- Domino tilings of Aztec diamond were introduced by Elkies-Kuperberg-Larsen-Propp'92. They proved that the number of tilings is equal to $2^{n(n+1)/2}$.
- **Exercise:** Check that Aztec diamond of size 2 has 8 domino tilings.
- Question: What happens when we consider a uniformly random domino tiling of a large Aztec diamond ?

Let us consider a chessboard coloring of Aztec diamond. It is useful to distinguish not two, but **four** different types of dominoes.



We will color these four types by different colors in the next picture.




We see that a uniformly random domino tiling has some structure !

Theorem (Jockush-Propp-Schor'98): Asymptotically a uniformly random tiling becomes frozen outside of a certain circle.

There are many more interesting properties of these tilings.


Let us consider dominoes of two types only:



Lemma: Each domino tiling is uniquely determined by dominoes of two types.

Idea of proof: Moving from north-west to south-east, we see that all missing dominoes can be reconstructed in a unique way.

(Johansson'02, '05) Let us put particles of two different types:



On level 0 we have 0 particles, then on levels 1 and 2 we have 1 particle, on levels 3 and 4 we have 2 particles, on levels 5 and 6 we have three particles.

Now we want to understand when a particle system corresponds to a domino tiling of the Aztec diamond.


A non-increasing sequence of integers $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_N$ is called a *signature* of length N .

We say that a signature λ of length N and signature μ of length $N - 1$ *interlace* if

$$\lambda_1 \geq \mu_1 \geq \lambda_2 \geq \mu_2 \geq \dots \geq \mu_{N-1} \geq \lambda_N.$$

Notation $\mu \prec \lambda$. Example: $(3, 3, 2) \prec (6, 3, 2, 0)$.


Geometrically, they differ by a *horizontal strip* (no two boxes in the same column).



We say that signatures λ and μ of length N differ by a *vertical strip* if $\lambda_i - \mu_i = \{0, 1\}$, for any $1 \leq i \leq N$.

Notation $\mu \prec_v \lambda$.

For example, $(3, 3, 1, 0, 0, 0) \prec_v (3, 3, 2, 1, 1, 1)$




Note that this example is obtained by symmetry with respect to $x = y$ from the previous one. This is a *transposition* of signatures / Young diagrams.

If λ is a signature of length N , then $\{\lambda_i + N - i\}_{i=1}^N$ is a collection of distinct integers. In the opposite direction, given a collection of distinct integers, one can construct a signature.

Consider all collections of signatures with nonnegative coordinates $\lambda^1, \dots, \lambda^N, \mu^1, \dots, \mu^{N-1}$, such that λ^i, μ^i have length i and

$$\emptyset \prec \lambda^1 \succ_v \mu^1 \prec \lambda^2 \succ_v \mu^2 \prec \dots \prec \lambda^N \succ_v (0, 0, \dots, 0)$$

Proposition. The set of collection of signatures is in bijection with domino tilings of Aztec diamond.



On each level boxes are numbered starting from 0. Distinct particles are in positions:

$$1; 1; (1, 3); (0, 2); (0, 1, 3); (0, 1, 2);$$

Making shifts, we obtain signatures:

$$\emptyset \prec 1 \succ_{\vee} 1 \prec (1, 2) \succ_{\vee} (0, 1) \prec (0, 0, 1) \succ_{\vee} (0, 0, 0)$$

Exercise: do this construction for a couple of domino tilings of Aztec diamond of size 3.

Condition:

$$\emptyset \prec \lambda^1 \succ_v \mu^1 \prec \lambda^2 \succ_v \mu^2 \prec \dots \prec \lambda^N \succ_v (0, 0, \dots, 0)$$

Proposition. The set of collections of signatures $(\lambda^1, \mu^1, \dots, \lambda^N)$ is in bijection with domino tilings of Aztec diamond.

Idea of proof: They correspond to black and white particles. Moving from south-east to north-west, one can check that the conditions on dominoes and on signatures coincide.

— How to compute the number of signatures with such conditions ?

— How to analyze the properties of uniformly random collection of signatures ?

Tools: Symmetric functions. Schur processes.